


Monitoring OA, HABs, and more from NANOOS Cha'ba and NEMO buoys

Jan Newton^{1, 4, 6} with many partners:

John Mickett¹, Simone Alin², Adrienne Sutton², Dick Feely²,
Stephanie Moore³, Terrie Klinger⁴, Nina Bednarsek⁵,
Parker MacCready⁶, Samantha Siedlecki⁷

1: UW-APL; 2: NOAA PMEL; 3: NOAA NWFSC; 4: UW-SMEA; 5: IPCC;
6: UW-Oceanography; 7: UW-JISAO





complex area...

2009 Science Issues

- WA coast is under sampled, physical dynamics are poorly resolved
- WA coast has seasonal hypoxia, strong inter-annual variation, but dynamics are different than off OR
- WA coast has a harmful algal bloom (HAB) “hot-spot” at Juan de Fuca eddy
- WA coast is exhibiting effects from OA now
- Current model accuracy is limited by data input

“A multi-platform high-resolution coastal ocean observing sensor array for researching Washington coastal waters and ecosystem response to climate change.”

Funded by Murdock Charitable Trust & UW now sustained as part of NANOOS

45" syntactic foam float
float depth 15 m in winter
10 m in summer

SBE 37 MicroCat

18 m (13 m in summer) stopper



(4) Benthos glass floats

Benthos 865 release
with recovery line canister





‘Like putting headlights on a car’

Pacific oysters gain from IOOS® data

About six years ago, production at some Pacific Northwest oyster hatcheries began declining at an alarming rate, posing severe economic impact and challenging a way of life held by shellfish growers for more than 130 years.

By 2008, the oyster harvest at Whiskey Creek, a major Oregon supplier to the majority of West Coast oyster farmers, plummeted 80 percent. At about the same time, corrosive, acidified seawater was hitting the shores of the Pacific.

Something had to be done. Oyster production accounts for more than \$84 million of the West Coast shellfish industry, which supports more than 3,000 jobs.

“When you see oyster shells dissolving in water, there’s a compelling need to know why,” says Bill Dewey of Taylor Shellfish Farms in Washington state.

Thanks to a \$500,000 federal investment in monitoring coastal seawater strengthened by data and observational information from the U.S. **Integrated Ocean Observing System (IOOS®)** and the **NOAA Ocean Acidification Program**, oyster hatcheries on the verge of collapse just a few years ago are again major contributors to the \$111 million West Coast shellfish industry.

IOOS is a NOAA-led interagency and regional effort aimed at “knowing” — that



IOOS partners in the Northwest Association of Networked Ocean Observing Systems (NANOOS) deployed this buoy in 2010 as part of a three-piece observing array to assess issues in the Northwest, including **ocean acidification**, **hypoxia and harmful algal blooms**, and **climate change**. The coastal buoy will aid computer models that predict ocean and atmospheric conditions. Known as “Chá bá,” the buoy is named for the Native American word (pronounced “chay buh”) for “whale tail.”

(Photo courtesy of Dr. John Payne, Pacific Ocean Shelf

Promoting
Economic Vitality


“Putting an IOOS buoy in the water is like putting headlights on a car. It lets us see changing water conditions in real time,” says Mark Wiegardt, co-owner of Whiskey Creek Shellfish Hatchery.








Realtime ChaBa Data: Temperature [°C]





ChaBa during cruise



UPWELLING WINDS


DOWNWELLING WINDS

wind direction (deg. T)




NEMO: Recent Successes

Measuring the “Blob” as it came ashore in the fall of 2014





- These were among the few full-depth, high-resolution measurements of this process.
- The measurements greatly complimented satellite observations of SST.


The “blob,” aka marine heat wave, is back....(or actually never really left)




Better definition of Columbia River influence




- PSEMP Marine Waters Workgroup. 2015. Puget Sound marine waters: 2014 overview. S. K. Moore, R. Wold, K. Stark, J. Bos, P. Williams, K. Dzinbal, C. Krembs and J. Newton (Eds). URL: <http://www.psp.wa.gov/PSEMP/PSmarinewatersoverview.php>.




A. Temperature at Cha'Ba at 85m



B. Salinity at Cha'Ba at 85 m




C. Dissolved oxygen (DO) at Cha'Ba at 85 m



ESP deployed on existing moored observatory in transport pathway


16



[Modified from Hickey et al. 2013]


NEMO Environmental Sample Processor (ESP)

- The ESP: a real-time harmful algal bloom (HAB)-detection system (MBARI)
- A complex lab-in-a-can that does in-situ DNA analysis, ID'ing target species and associated toxin
- WA razor clam fishery alone experiences an estimated \$24.4 million in lost revenue when closed to HABs. Dungeness crab also important.
- Significant human health aspect
- Can cause severe ecosystem impacts on higher trophic levels.
- The real-time mooring will be an early-detection system to enable rapid resource management and human health decisions.
- Will be integrated into the NEMO subsurface mooring, with first operational deployment scheduled for this May.




NEMO Environmental Sample Processor

- APL lead of 7-institution effort.
- designed and built the mooring infrastructure, pressure case, power system, telemetry system, and pump sampling system.
- significantly expands the scope of deployment environments for the ESP.





ITEM NO.	DESCRIPTION	QTY.
1	FRAME BASE, GALVANIZED STEEL	1
2	NEMO MOORING CONTROLLER	1
3	ESP BATTERY HOUSING	2
4	ESP ASSEMBLY	1
5	750M SYNTACTIC FOAM	1
6	THRU LIFT FRAME, 316SS	1
7	TELEDYNE SENTINEL ADCP	1
8	FIBERGLASS GRATING	2
9	SATLANTIC SUNA	1
10	SEABIRD 37	1
11	SATLANTIC SEA FET	1



data transmitted to surface mooring
by cell modem in telebuoy
(~400 m from surface mooring)

data telemetered
to shore near real-time





Cha'ba is a national OA buoy...



PMEL
CARBON PROGRAM

Search

CONTACT US

PARTNERS

INTERNAL

SITE MAP

Home

Research

Observations

Outreach & Education

Data Portal

People

News



Strategy for OA Observations

PMEL is developing a global network of ocean acidification observations

The existing global carbon observatory network of repeat hydrographic surveys, time-series stations and ship-based underway surface observations in the open ocean provide a strong foundation of carbon chemistry observations to begin addressing the problem of ocean acidification. Indeed, much of our present understanding of the long-term changes in the carbon system is derived from the repeat ocean sections and time-series measurements.

A major project for our group is to **expand the global moored and ship-based network by adding pH and other biogeochemical measurements** to provide important information on the changing conditions in the open ocean and coastal waters. See the map of planned monitoring sites to the right. This network will provide a better understanding of the temporal and spatial scales of variability in ocean carbon chemistry and biology and the observational basis for developing predictive models for future changes in ocean acidification and its consequences for marine ecosystems.



Location of planned OA monitoring and research sites and affiliated NOAA labs.

RELATED STORIES




Ocean Acidifica ...
Fundamental changes in seawater chemistry are occurring throu ...




NOAA OA Plan
NOAA researchers and managers are working to coordinate ocean ...



U.S. **coastal and estuarine environments** do not currently have coordinated carbon observing networks, as in the open ocean, and are presently grossly under-sampled. There is a critical need for intensive time series measurements on moored buoys and repeat hydrographic cruises in high productivity coastal and estuarine systems as CO₂ and carbonate ion concentrations in these waters can vary substantially on timescales from hours to decades due to tides, photosynthesis, and river or ground water inputs. In response to that need, we are adding carbon and pH sensors to




Wind



pH




DO and pH relations



2015 Science Reality

- WA coast is sampled in near real-time; physical dynamics revealed and published
- WA coast has seasonal hypoxia; dynamics can be observed by tribes and agencies in real-time
- WA coast has a HAB “hot-spot”; now status will be monitored for toxicity in real-time
- WA coast is exhibiting effects from OA now, observed by a NOAA OAP buoy (Cha’ba) year-round
- Current model accuracy is increased and skill can be verified by the on-line “comparator”

NANOOS observations and data



Washington Ocean Acidification Center implementing key recommendations



Coordinate and synthesize science to:

- 1. Assess water conditions and what's driving ocean acidification***
 - Monitoring (both in natural environment and at shellfish hatcheries)
- 2. Provide forecasts to facilitate adaptation***
 - Forecast modeling
- 3. Assess how local species respond***
 - Biological experiments
- 4. Inform aquaculture practices***
 - Shellfish culture adaptation

1. Assess OA and biological response


Strategies:

- Both **chemistry** (DIC, TA) and **biology** measurements
- **Temporal** trends (buoys) & **spatial** coverage (surveys)
- **Leverages** existing networks




Washington water chemistry

$$\Omega_{\text{arag}}$$





Pteropods as bio-indicators




No Dissolution






Type I: Moderate



Type II: Severe




Type III: Severe




WA OA Center monitoring: *bioindicator*

- Pteropod shells show signs of dissolution
- Patterns in time and space help us understand impacts and drivers




La Push, coastal WA




Whidbey basin, Puget Sound

Near-term predictions of OA in Washington

3-Day forecast appears daily on NANOOS




Breaking ground to seasonal ecosystem forecasting: hypoxia, OA, sardines




The GOA-ON interactive data portal

Featuring global OA data, asset inventory, metadata, data synthesis products, etc.




We are GOA-ON !





Source: IPCC Report



**OA is
a global condition
with
local effects**

